

HADRONIZATION IN $\tau \rightarrow KK\pi\nu_\tau$ DECAYS

P. ROIG

*Instituto de Física Corpuscular, IFIC, CSIC-Universitat de València.
Apt. de Correus 22085, E-46071 València, Spain*



Hadronization in $\tau \rightarrow KK\pi\nu_\tau$ decays is driven by both vector and axial-vector currents that we study, guided by the following principles: The $1/N_C$ expansion -worked out at leading order, considering only the contribution of the lightest spin one resonances-, approximate chiral symmetry at low energies and the appropriate asymptotic behaviour we demand to the associated form factors. All these features are implemented in the resonance theory. Most of its couplings are determined by imposing the short-distance requirements of vector and axial-vector spectral functions within QCD. We plan to improve our prediction of the hadronic spectra using recently available experimental data.

1 Introduction

The mass of the τ lepton, ~ 1.8 GeV, makes it privileged for it connects the hadronic and leptonic worlds through its semileptonic decays in the cleanest possible way. On the other hand, it sets the typical energy scale of these processes in a region where the fundamental theory of strong interactions, QCD¹, becomes computationally useless.

One may, of course, rely on parameterizations of the assumed dynamics that have proved to be very successful in the modes involving two and three pions, namely the Kühn-Santamaría (KS) model², but it would be desirable to explain the experimental data within a framework that resembles QCD as much as possible.

On the very low energy domain χ PT^{3,4} is the QCD dual at energies well below the $\rho(770)$ mass, i.e., $E \ll 0.6$ GeV. As expected, however, χ PT only applies to a small part of the phase space in τ decays⁵. The involved hadronization can not be understood without including explicitly the resonances that are exchanged in the process. The most common procedure is to rely on

the phenomenologically successful notion of vector meson dominance⁶, as we do.

The easiest way to incorporate the resonances is through Breit-Wigner functions, including phase-space motivated off-shell widths and a weighted interplay between the fundamental and excited resonances as in Ref.². This has been shown^{7,8} to violate chiral symmetry at NLO in the chiral expansion. In addition, the assumed structure of exchanged resonances in the $KK\pi$ modes⁹ does not cover all those allowed, as can be checked by comparison with Ref.¹⁰. Furthermore, Ref.⁹ includes different mass and width for the $\rho(770)$ depending on its appearance on the axial-vector current or the vector one. Finally, two multiplets of vector resonances are used⁹ in the axial-vector current, while three of them are employed in the vector one. The aim of our work is to achieve a description of the considered decays closer to QCD than this one. CLEO¹¹, BaBar¹² and Belle¹³ have been collecting good quality data on $\tau \rightarrow KK\pi\nu_\tau$ decays. CLEO¹¹ announced that the parameterization in Ref.⁹ was unable to fit their data so that they reshaped it violating the Wess-Zumino normalization emanating from the chiral anomaly of QCD, as was put forward in Ref.¹⁴. BaBar has not published these results yet. However, they have managed¹⁵ to split with great precision the isoscalar and isovector contributions in $e^+e^- \rightarrow KK\pi$ so that, under the isospin symmetry we can use it for the considered τ decays, as it has been done in Ref.¹⁶.

The fact that CLEO and BaBar disagree with respect to their respective results being explained or not by a model generalizing KS², on the one hand, and the inconsistencies in the treatment followed in Refs.^{9,17} -as stated in the previous paragraph- are the motivations for our work trying to explain the experimental measurements while learning as much as possible about the hadronization of the involved QCD currents.

2 THE RESONANCE CHIRAL THEORY OF QCD

As opposed to the very low energy sector, the effective theory of QCD at intermediate energies remains unknown. Our aim is to resemble the underlying theory as much as possible. For this, we rely on Resonance Chiral Theory (R χ T), as introduced in the article¹⁸, which is built upon the approximate chiral symmetry of low-energy QCD for the lightest pseudoscalar mesons and unitary symmetry for the resonances. Weinberg's Theorem^{3,19} ensures the correctness of writing a phenomenological Lagrangian in terms of mesons fulfilling the basic symmetries of light-flavoured QCD. Its large- N_C limit²⁰ guides the perturbative expansion of R χ T. At LO in $1/N_C$, meson dynamics is described by tree level diagrams obtained from an effective local Lagrangian including the interactions among an infinite number of stable resonances. In τ decays we need to include finite resonance widths (a NLO effect in this expansion) as we do within our framework²¹. We are also departing from the $N_C \rightarrow \infty$ limit because we consider just one multiplet of resonances per set of quantum numbers (single resonance approximation²²) and not the infinite tower predicted that cannot be included in a model independent way. Representing the resonances in the antisymmetric tensor formalism proves to be more convenient²³ in processes involving mesons as asymptotic states, instead of the Proca²⁴ or mixed²⁵ ones. The relevant part of the R χ T Lagrangian is^{18,26,27,28}:

$$\begin{aligned} \mathcal{L}_{R\chi T} = & \frac{F^2}{4} \langle u_\mu u^\mu + \chi_+ \rangle + \frac{F_V}{2\sqrt{2}} \langle V_{\mu\nu} f_+^{\mu\nu} \rangle + \frac{iG_V}{\sqrt{2}} \langle V_{\mu\nu} u^\mu u^\nu \rangle + \frac{F_A}{2\sqrt{2}} \langle A_{\mu\nu} f_-^{\mu\nu} \rangle \\ & + \mathcal{L}_{\text{kin}}^V + \mathcal{L}_{\text{kin}}^A + \sum_{i=1}^5 \lambda_i \mathcal{O}_{VAP}^i + \sum_{i=1}^7 \frac{c_i}{M_V} \mathcal{O}_{VJP}^i + \sum_{i=1}^4 d_i \mathcal{O}_{VVP}^i + \sum_{i=1}^5 \frac{g_i}{M_V} \mathcal{O}_{VPPP}^i, \end{aligned} \quad (1)$$

where all couplings are real, being F the pion decay constant in the chiral limit. The notation is that of Ref.¹⁸. Here and in the following P stands for the lightest pseudoscalar mesons and

A and V for the (axial-)vector mesons. Furthermore, all couplings in the second line are defined to be dimensionless. For the explicit form of the operators in the last line, see^{26, 27, 28}.

In order to inherit the maximum possible features of QCD, to be implemented in R χ T, we still have to exploit the matching between order parameters of spontaneous chiral symmetry breaking with partonic QCD related quantities. The matching of n -point Green Functions in the OPE of QCD and in R χ T has been shown to be a fruitful procedure^{23, 26, 27, 29, 30, 31}. Additionally, we will demand to the vector and axial-vector form factors a Brodsky-Lepage-like behaviour³². While symmetry fully determines the structure of the operators, it is the QCD-ruled short-distance behaviour who restricts certain combinations of couplings rendering R χ T predictive provided we fix these few remaining parameters restoring to phenomenology: F_V could be extracted from the measured $\Gamma(\rho^0 \rightarrow e^+e^-)$, G_V from $\Gamma(\rho^0 \rightarrow \pi^+\pi^-)$, F_A from $\Gamma(a_1 \rightarrow \pi\gamma)$ and the λ_i 's from $\Gamma(a_1 \rightarrow \rho\pi)$ which starts on the $\tau \rightarrow 3\pi\nu_\tau$ processes themselves. $\Gamma(\omega \rightarrow \pi\gamma)$, $\Gamma(\omega \rightarrow 3\pi)$ and the $\mathcal{O}(p^6)$ correction to $\Gamma(\pi \rightarrow \gamma\gamma)$ may give us information on the remaining couplings²⁷.

3 FORM FACTORS IN $\tau^- \rightarrow (KK\pi)^- \nu_\tau$

The decay amplitudes for all charge channels can be written as

$$\mathcal{M} = -\frac{G_F}{\sqrt{2}} V_{ud} \bar{u}_{\nu_\tau} \gamma^\mu (1 - \gamma_5) u_\tau \mathcal{H}_\mu, \quad (2)$$

where

$$\begin{aligned} \mathcal{H}_\mu &= \langle P(p_1)P(p_2)P(p_3) | (\mathcal{V}_\mu - \mathcal{A}_\mu) e^{i\mathcal{L}_{QCD}} | 0 \rangle = \\ &V_{1\mu} F_1^A(Q^2, s_1, s_2) + V_{2\mu} F_2^A(Q^2, s_1, s_2) + Q_\mu F_3^A(Q^2, s_1, s_2) + iV_{3\mu} F_4^V(Q^2, s_1, s_2), \end{aligned} \quad (3)$$

and

$$\begin{aligned} V_{1\mu} &= \left(g_{\mu\nu} - \frac{Q_\mu Q_\nu}{Q^2} \right) (p_2 - p_1)^\nu, \quad V_{2\mu} = \left(g_{\mu\nu} - \frac{Q_\mu Q_\nu}{Q^2} \right) (p_3 - p_1)^\nu, \\ V_{3\mu} &= \varepsilon_{\mu\nu\rho\sigma} p_1^\nu p_2^\rho p_3^\sigma, \quad Q_\mu = (p_1 + p_2 + p_3)_\mu, \quad s_i = (Q - p_i)^2. \end{aligned} \quad (4)$$

Here F_i , $i = 1, 2, 3$ correspond to the axial-vector current while F_4 drives the vector current. The form factors F_1 and F_2 have a transverse structure in the total hadron momenta, Q^μ , and drive a $J^P = 1^+$ transition. The scalar form factor, F_3 , vanishes with the mass of the Goldstone bosons (chiral limit) and, accordingly, gives a tiny contribution. In Ref.⁸ the vector form factor was not contributing, so we do not only intend to confirm or refuse the bounds on the $\{\lambda_i\}_{i=1}^5$, but to explore the vector current sector of the resonance Lagrangian. The explicit expressions of the form factors can be found in Refs.^{28, 10}.

4 ASYMPTOTIC BEHAVIOUR AND QCD CONSTRAINTS

Computing the Feynman diagrams involved shows that the result depends only on three combinations of the $\{\lambda_i\}_{i=1}^5$, four of the $\{c_i\}_{i=1}^7$, two of the $\{d_i\}_{i=1}^4$ and four of the $\{g_i\}_{i=1}^5$. The number of free parameters has been reduced from 24 to 15.

We require the form factors of the A^μ and V^μ currents into $KK\pi$ modes vanish at infinite transfer of momentum. As a result, we obtain constraints³³ among all axial-current couplings but λ_0 , that are also the most general ones satisfying the demanded asymptotic behaviour in $\tau \rightarrow 3\pi\nu_\tau$. Proceeding analogously with the vector current form factor results in five additional restrictions³³. From the 24 initially free couplings in Eq. (1), only five remain free: c_4 ,

$c_1 + c_2 + 8c_3 - c_5$, $d_1 + 8d_2 - d_3$, g_4 and g_5 . After fitting $\Gamma(\omega \rightarrow 3\pi)$ -using some relations obtained in²⁷- only c_4 and g_4 remain unknown.

Employing isospin symmetry³⁶ we are able to provide a theoretical expression for $\sigma(e^+e^- \rightarrow KK\pi)$ that we have fitted to BaBar data¹⁵ obtaining $c_4 = -0.052 \pm 0.003$ and $g_4 = -0.20^{+0.08}_{-0.12}$. Remarkably, this procedure has allowed us to be sensitive to the sign of c_4 improving the work done in³³. The results are represented in Fig 1.




Figure 1: Fit of our computation for $\sigma(e^+e^- \rightarrow KK\pi)$ to BaBar data. At higher energies than those shown the contribution of excited resonances we do not have taken into account becomes relevant.

Acknowledgments

I wish to thank Jean Tran Thanh Van for the excellent organization, and the warm and cosy atmosphere of the QCD and High Energy Interactions 2008 meeting in La Thuile (Italy). I acknowledge B. Malaescu for pointing out to me that BaBar $e^+e^- \rightarrow KK\pi$ data was already available. This work has been done in collaboration with D. Gómez-Dumm, A. Pich and J. Portolés. I wish to thank J. P. for a careful revision of the manuscript. P. R. is supported by a FPU contract (MEC). This work has been supported in part by the EU MRTN-CT-2006-035482 (FLAVIANet), by MEC (Spain) under grant FPA2007-60323, by the Spanish Consolider-Ingenio 2010 Program CPAN (CSD2007-00042) and by Generalitat Valenciana under grant GVACOMP2007-156.

References

1. H. Fritzsch, M. Gell-Mann and H. Leutwyler, *Phys. Lett. B* **47**, 365 (1973).
2. J. H. Kühn and A. Santamaría, *Z. Phys. C* **48**, 445 (1990).
3. S. Weinberg, *Physica A* **96**, 327 (1979).
4. J. Gasser and H. Leutwyler, *Annals of Phys.* **158**, 142 (1984), *Nucl. Phys. B* **250**, 465 (1985).
5. G. Colangelo, M. Finkemeier and R. Urech, *Phys. Rev. D* **54**, 4403 (1996).
6. J. H. Kühn and F. Wagner, *Nucl. Phys. B* **236**, 16 (1984). A. Pich, in Proceedings of the *Tau Charm Factory Workshop* Ed. L. V. Beers, SLAC (1989).
7. J. Portolés, *Nucl. Phys. Proc. Suppl.* **98** 210 (2001).
8. D. Gómez Dumm, A. Pich and J. Portolés, *Phys. Rev. D* **69**, 073002 (2004).
9. M. Finkemeier and E. Mirkes, *Z. Phys. C* **69**, 243 (1996).
10. D. Gómez-Dumm, P. Roig, A. Pich, J. Portolés, work in preparation.
11. F. Liu [CLEO Collaboration], *Nucl. Phys. Proc. Suppl.* **123**, 66 (2003). T. E. Coan *et al.* [CLEO Collaboration], *Phys. Rev. Lett.* **92**, 232001 (2004).
12. I. M. Nugent [BABAR Collaboration], *Nucl. Phys. Proc. Suppl.* **169**, 98 (2007).
13. [Belle Collaboration], work in preparation (S. Eidelman, private communication).
14. J. Portolés, *Nucl. Phys. Proc. Suppl.* **144**, 3 (2005).
15. B. Aubert *et al.* [BABAR Collaboration], *Phys. Rev. D* **77** (2008) 092002.
16. M. Davier, S. Descotes-Genon, A. Hocker, B. Malaescu, Z. Zhang, *Eur. Phys. J. C* **56**, 305 (2008).
17. M. Finkemeier, J. H. Kühn and E. Mirkes, *Nucl. Phys. Proc. Suppl.* **55C**, 169 (1997).
18. G. Ecker, J. Gasser, A. Pich and E. de Rafael, *Nucl. Phys. B* **321**, 311 (1989).
19. H. Leutwyler, *Annals of Phys.* **235**, 165 (1994).
20. G. 't Hooft, *Nucl. Phys. B* **72**, 461 (1974), 75 461 1974. E. Witten, *Nucl. Phys. B* **160**, 57 (1979).
21. D. Gómez Dumm, A. Pich and J. Portolés, *Phys. Rev. D* **62**, 054014 (2000).
22. M. Knecht, S. Peris, M. Perrottet and E. de Rafael, *Phys. Rev. Lett.* **83**, 5230 (1999), *Jour. of High Energy Phys.* **0211**, 003 (2002).
23. G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. de Rafael, *Phys. Lett. B* **223**, 425 (1989).
24. J. F. Donoghue, C. Ramírez, G. Valencia *Phys. Rev. D* **39**, 1947 (1989).
25. K. Kampf, J. Novotny, J. Trnka *Eur. Phys. J. C* **50**, 385 (2007).
26. V. Cirigliano, G. Ecker, M. Eidemüller, A. Pich and J. Portolés, *Phys. Lett. B* **596**, 96 (2004).
27. P. D. Ruiz-Femenía, A. Pich and J. Portolés, *Jour. of High Energy Phys.* **0307**, 003 (2003).
28. P. Roig, Treball d'Investigació, Universitat de València, 2006.
29. G. Amorós, S. Noguera and J. Portolés, *Eur. Phys. J. C* **27**, 243 (2003).
30. M. Knecht and A. Nyffeler, *Eur. Phys. J. C* **21**, 659 (2001).
31. V. Cirigliano, G. Ecker, M. Eidemüller, R. Kaiser, A. Pich and J. Portolés, *Nucl. Phys. B* **753**, 139 (2006).
32. G. P. Lepage and S. J. Brodsky, *Phys. Rev. D* **22**, 2157 (1980).
33. P. Roig, *AIP Conf. Proc.* **964**, 40 (2007).
34. W.-M. Yao *et al.*, *Journal of Physics G* **33**, 1 (2006).
35. J. J. Gómez-Cadenas, M. C. González-García and A. Pich, *Phys. Rev. D* **42**, 3093 (1990).
36. R. Barate *et al.* [The ALEPH Collaboration] *Eur. Phys. Jour. C* **11**, 599 (1999).